



## The Iterative Solution of Taylor Formula for Partial Differential Equation

Yang GAO

Independent Researcher, China

**Citation:** Yang GAO (2026) The Iterative Solution of Taylor Formula for Partial Differential Equation. *J. of Mod Phy & Quant Neuroscience* 2(2), 1-07. WMJ/JPQN-154

### Abstract

This paper discusses the relation between Taylor's formula and partial differential equation. Taylor formula iteration method can resolve partial differential equation.  $u(x,t)$  be expanded at  $t = 0$  or  $t = 1$  by Taylor formula. Coefficient of Taylor formula  $u_t(x,0), u_{tt}(x,0) \dots$  can be expressed by partial differential equation. The method can solve nonlinear differential equation. Generalized Taylor's formula can solve fractional partial differential equation. The method is very important way that resolving partial differential equation. The method also can resolve those equations from [1-9].

This article refers to the literature [10]. Taylor formula iteration method belongs to logical thinking.

\*Corresponding author: Yang GAO, Independent Researcher, China.

Submitted: 11.03.2026

Accepted: 17.03.2026

Published: 31.03.2026

**Keyword:** Taylor's Formula, Iteration Method, Nonlinear, Partial Differential Equation

### Introduction

This paper introduce that Taylor formula iteration method resolve partial differential equation. In this paper, six examples are used to introduce Taylor formula iteration method to solve partial differential equation. This paper also introduces that Generalized Taylor's formula can solve fractional partial differential equation. The iterative method of Taylor formula is an important and useful method to solve partial differential equation. The solution of Taylor formula iteration method belongs to  $C^\infty$ .

Variable Coefficient Problem

We consider equation as following:

$$xu_{tt}(x,t) - (xu_x(x,t))_x = 0, \quad (1)$$

$$u(x,0) = x^2, \quad (2)$$

$$u_t(x,0) = 0. \quad (3)$$

We solve (1) by Taylor formula iteration method as following:

$$\begin{aligned}
 (4) \quad & xu_{tt}(x, t) - (xu_x(x, t))_x = 0, \\
 (5) \quad & u_{tt} = u_{xx} + \frac{1}{x}u_x, \\
 (6) \quad & u_{tt}(x, 0) = u_{xx}(x, 0) + \frac{1}{x}u_x(x, 0), \\
 (7) \quad & u_{xx}(x, 0) = (x^2)''_{xx} = 2, \\
 (8) \quad & u_x(x, 0) = (x^2)'_x = 2x, \\
 (9) \quad & u_{tt}(x, 0) = 2 + 2 = 4, \\
 (10) \quad & u_{ttt} = u_{txx} + \frac{1}{x}u_{tx}, \\
 (11) \quad & u_{ttt}(x, 0) = u_{txx}(x, 0) + \frac{1}{x}u_{tx}(x, 0), \\
 (12) \quad & u_t(x, 0) = 0, \\
 (13) \quad & u_{txx}(x, 0) = 0, \\
 (14) \quad & u_{tx}(x, 0) = 0, \\
 (15) \quad & u_{ttt}(x, 0) = 0,
 \end{aligned}$$

And we have:

$$u_{ttt}(x, 0) = u_{ttt}(x, 0) = \dots = 0 \tag{16}$$

By Taylor’s formula, we get as following:

$$u(x, t) = u(x, 0) + u_t(x, 0)t + u_{tt}(x, 0)\frac{t^2}{2!} + u_{ttt}(x, 0)\frac{t^3}{3!} + \dots, \tag{17}$$

$$u(x, t) = x^2 + 2t^2. \tag{18}$$

Solution of equation (1) , $u(x, t) = x^2 + 2t^2$ .

### Two-Dimensional Heat Conduction Equation Solution

We study the equation as following:

$$u_t - tx(u_{xx} + u_{yy}) = t^2, \tag{19}$$

$$u(x, y, 0) = xy + y^3. \tag{20}$$

Next, we solve (19) by Taylor formula iteration method,

$$u_t = tx(u_{xx} + u_{yy}) + t^2, \tag{21}$$

Let  $t = 0$  on (21),

$$u_t(x, y, 0) = 0, \tag{22}$$

On equation (21), finding 1-order partial derivative of  $t$  on both sides,

We have:

$$utt = x(uxx + uyy) + tx(utxx + utyy) + 2t, \tag{23}$$

$$u_{xx}(x, y, 0) = 0, \tag{24}$$

$$u_{yy}(x,y,0) = 6y, \tag{25}$$

Next,  $t = 0$  on (23),

$$u_u(x,y,0) = 6xy, \tag{26}$$

$$uttt = 2x(utxx + utyy) + tx(uttxx + uttyy) + 2, \tag{27}$$

$$u_{ttt}(x,y,0) = 0 = (6xy)_{xx} = 0 \tag{28}$$

$$u_{tt}(x,y,0) = 0, \tag{29}$$

$$u_{tt}(x,y,0) = 0, \tag{30}$$

Next,  $t = 0$  on (27),

$$u_{tt}(x,y,0) = 2, \tag{31}$$

$$utttt = 3x(uttxx + uttyy) + tx(utttxx + utttyy), \tag{32}$$

$$u_{tttt}(x,y,0) = 0, \tag{33}$$

So we have : (34)

$$u_{tttt}(x,y,0) = u_{tttt}(x,y,0) = \dots = 0, \tag{35}$$

By Taylor’s formula ,we get as following:

$$u(x, y, t) = u(x, y, 0) + u_t(x, y, 0)t + u_{tt}(x, y, 0)\frac{t^2}{2!} + u_{ttt}(x, y, 0)\frac{t^3}{3!} + \dots, \tag{36}$$

$$u(x, y, t) = (3t^2 + 1)xy + y^3 + \frac{t^3}{3}. \tag{37}$$

Solution of (19),  $u(x, y, t) = (3t^2 + 1)xy + y^3 + \frac{t^3}{3}$ .

### The Third Problem With Boundary Values

We consider following equation:

$$ut - 4uxx = \text{cost}, \tag{38}$$

$$u(x,0) = \text{cos}x, \tag{39}$$

$$u_x(0,t) = u_x(1,t) = 0. \tag{40}$$

By Taylor formula iteration method,we have:

$$u_t = 4u_{xx} + \text{cost}, \tag{41}$$

$$u_t(x,0) = 4u_{xx}(x,0) + 1, \tag{42}$$

$$u_t(x,0) = -4\text{cos}x + 1, \tag{43}$$

$$utt = 4u_{txx} - \text{sint}, \tag{44}$$

$$u_{tt}(x,0) = 4u_{txx}(x,0), \tag{45}$$

$$u_{tt}(x,0) = 4^2 \text{cos}x, \tag{46}$$

$$uttt = 4u_{ttxx} - \text{cost}, \tag{47}$$

$$u_{ttt}(x,0) = 4u_{ttxx}(x,0) - 1, \tag{48}$$

$$u_{ttt}(x,0) = -4^3 \cos x - 1, \tag{49}$$

$$utttt = 4u_{tttx} + \sin t, \tag{50}$$

$$u_{ttt}(x,0) = 4u_{tttx}(x,0), \tag{51}$$

$$u_{ttt}(x,0) = 4^4 \cos x, \tag{52}$$

$$uttttt = 4u_{ttttx} + \cos t, \tag{53}$$

$$u_{ttt}(x,0) = -4^5 \cos x + 1, \tag{54}$$

$$\tag{55}$$

By Taylor’s formula ,we get as following:

$$u(x, t) = u(x, 0) + u_t(x, 0)t + u_{tt}(x, 0)\frac{t^2}{2!} + u_{ttt}(x, 0)\frac{t^3}{3!} + \dots, \tag{56}$$

$$u(x, t) = \sum_{n=0}^{\infty} \frac{(-4)^n}{n!} t^n \cos x + \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n + 1)!} t^{2n+1}, \tag{57}$$

$$\sin t = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n + 1)!} t^{2n+1}, \tag{58}$$

We have:

$$u(x, t) = e^{-4t} \cos x + \sin t. \tag{59}$$

We take the best of Fourier expansion:

$$a_n(t) = 2 \int_0^1 (e^{-4t} \cos x) \cos(n\pi x) dx, \tag{60}$$

We get the solution of (38):

$$u(x, t) = \sum_{n=1}^{\infty} a_n(t) \cos(n\pi x) + e^{-4t} \sin 1 + \sin t. \tag{61}$$

### Fractional Partial Differential Equation

We consider following fractional partial differential equation:

$$\frac{\partial^\alpha u(x, y, t)}{\partial t^\alpha} = D_x^\beta u(x, y, t) + D_y^\gamma u(x, y, t) + u(x, y, t) \tag{62}$$

$$u(x, y, 0) = q(x, y). \tag{63}$$

where  $q(x,y)$ is known integral polynomial.

The definition of Caputo fractional derivative about:

$q_1(x, y)$ is known function.

We consider  $\frac{\partial^{2\alpha} u(x, y, 0)}{\partial t^{2\alpha}}$ . On equation (62),finding  $\alpha$ -order partial derivative of  $t$  on both sides,

We have:

$$\frac{\partial^{2\alpha} u(x, y, t)}{\partial t^{2\alpha}} = D_x^\beta \frac{\partial^\alpha u(x, y, t)}{\partial t^\alpha} + D_y^\gamma \frac{\partial^\alpha u(x, y, t)}{\partial t^\alpha} + \frac{\partial^\alpha u(x, y, t)}{\partial t^\alpha}, \tag{71}$$

$$\frac{\partial^{2\alpha} u(x, y, 0)}{\partial t^{2\alpha}} = D_x^\beta \frac{\partial^\alpha u(x, y, 0)}{\partial t^\alpha} + D_y^\gamma \frac{\partial^\alpha u(x, y, 0)}{\partial t^\alpha} + \frac{\partial^\alpha u(x, y, 0)}{\partial t^\alpha}, \tag{72}$$

$$\frac{\partial^{2\alpha} u(x, y, 0)}{\partial t^{2\alpha}} = D_x^\beta q_1(x, y) + D_y^\gamma q_1(x, y) + q_1(x, y), \tag{73}$$

where

$$\frac{\partial^{2\alpha} u(x, y, 0)}{\partial t^{2\alpha}} = q_2(x, y), \tag{74}$$

$q_1(x,y)$  is known function.

We consider  $\frac{\partial^{2\alpha} u(x,y,0)}{\partial t^{2\alpha}}$ . On equation (62), finding  $\alpha$ -order partial derivative of  $t$  on both sides, We have:

$$\frac{\partial^{3\alpha} u(x,y,t)}{\partial t^{3\alpha}} = D_x^\beta \frac{\partial^{2\alpha} u(x,y,t)}{\partial t^{2\alpha}} + D_y^\gamma \frac{\partial^{2\alpha} u(x,y,t)}{\partial t^{2\alpha}} + \frac{\partial^{2\alpha} u(x,y,t)}{\partial t^{2\alpha}}, \quad (75)$$

$$\frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}} = D_x^\beta \frac{\partial^{2\alpha} u(x,y,0)}{\partial t^{2\alpha}} + D_y^\gamma \frac{\partial^{2\alpha} u(x,y,0)}{\partial t^{2\alpha}} + \frac{\partial^{2\alpha} u(x,y,0)}{\partial t^{2\alpha}}, \quad (76)$$

$$\frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}} = D_x^\beta q_2(x,y) + D_y^\gamma q_2(x,y) + q_2(x,y), \quad (77)$$

where

$$\frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}} = q_3(x,y), \quad (78)$$

$q_3(x,y)$  is known function. We consider  $\frac{\partial^{4\alpha} u(x,y,0)}{\partial t^{4\alpha}}$ . On equation (62), finding  $3\alpha$ -order partial derivative of  $t$  on both sides, we have:

$$\frac{\partial^{4\alpha} u(x,y,t)}{\partial t^{4\alpha}} = D_x^\beta \frac{\partial^{3\alpha} u(x,y,t)}{\partial t^{3\alpha}} + D_y^\gamma \frac{\partial^{3\alpha} u(x,y,t)}{\partial t^{3\alpha}} + \frac{\partial^{3\alpha} u(x,y,t)}{\partial t^{3\alpha}}, \quad (79)$$

$$\frac{\partial^{4\alpha} u(x,y,0)}{\partial t^{4\alpha}} = D_x^\beta \frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}} + D_y^\gamma \frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}} + \frac{\partial^{3\alpha} u(x,y,0)}{\partial t^{3\alpha}}, \quad (80)$$

$$\frac{\partial^{4\alpha} u(x,y,0)}{\partial t^{4\alpha}} = D_x^\beta q_3(x,y) + D_y^\gamma q_3(x,y) + q_3(x,y), \quad (81)$$

where

$$\frac{\partial^{4\alpha} u(x,y,0)}{\partial t^{4\alpha}} = q_4(x,y), \quad (82)$$

$q_4(x,y)$  is known function. We have:  $\frac{\partial^{n\alpha} u(x,y,0)}{\partial t^{n\alpha}} = q_n(x,y)$  (83)

$q_n(x,y)$  is known function.

By Generalized Taylor's formula:

$$u(x,y,t) = \sum_{j=0}^N \frac{t^{j\alpha}}{\Gamma(j\alpha + 1)} \frac{\partial^{j\alpha} u(x,y,0)}{\partial t^{j\alpha}} + \dots \quad (84)$$

So, we have the solution of (62):

$$u(x,y,t) = q(x,y) + q_1(x,y) \frac{t^\alpha}{\Gamma(\alpha + 1)} + q_2(x,y) \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} + \dots \quad (85)$$

### Nonlinear KdV Equation

We consider the wave equation as following:

$$u_t - 6uu_x + u_{xxx} = 0, \quad (86)$$

$$u(x,0) = x. \quad (87)$$

We have:

$$u_t = 6uu_x - u_{xxx}, \quad (88)$$

$$u_t(x,0) = 6u(x,0)u_x(x,0) - u_{xxx}(x,0), \quad (89)$$

$$u(x,0) = x, \quad (90)$$

$$u_x(x,0) = 1, \quad (91)$$

$$u_{xxx}(x,0) = 0, \quad (92)$$

$$u_t(x,0) = 6x, \quad (93)$$

On equation (88), finding 1-order partial derivative of t on both sides, we have:

$$u_{tt} = 6(u_t u_x + uu_{tx}) - u_{txxx}, \tag{94}$$

$$u_{tt}(x, 0) = 6(u_t(x, 0)u_x(x, 0) + u(x, 0)u_{tx}(x, 0)) - u_{txxx}(x, 0), \tag{95}$$

$$u_{tx}(x, 0) = 6, \tag{96}$$

$$u_{txxx}(x, 0) = 0, \tag{97}$$

$$u_{tt}(x, 0) = 2 \cdot 6^2 x, \tag{98}$$

On equation (94), finding 1-order partial derivative of t on both sides, we have:

$$u_{ttt} = 6(u_{tt}u_x + 2u_t u_{tx} + uu_{ttx}) - u_{ttxxx}, \tag{99}$$

$$u_{ttt}(x, 0) = 6(u_{tt}(x, 0)u_x(x, 0) + 2u_t(x, 0)u_{tx}(x, 0) + u(x, 0)u_{ttx}(x, 0)) - u_{ttxxx}(x, 0),$$

$$u_{tx}(x, 0) = 6, \tag{100}$$

$$u_{ttx}(x, 0) = 2 \cdot 6^2, \tag{101}$$

$$u_{ttt}(x, 0) = 6^4 x. \tag{102}$$

By Taylor’s formula ,we get as following:

$$u(x, t) = u(x, 0) + u_t(x, 0)t + u_{tt}(x, 0)\frac{t^2}{2!} + u_{ttt}(x, 0)\frac{t^3}{3!} + \dots, \tag{103}$$

We have:

$$u(x, t) = x + 6xt + 6^2 xt^2 + 6^3 xt^3 + 6^4 xt^4 + \dots, \tag{104}$$

$$\tag{105}$$

$$u(x, t) = \frac{x}{1 - 6t}. \tag{106}$$

So we can get the solution of (86),  $u(x, t) = \frac{x}{1 - 6t}$ .

### Nonlinear sine-Gordon Equation

We consider following constant coefficient equation

$$u_{tt} - c^2 u_{xx} + \alpha \sin u = 0, \tag{107}$$

$$u(x, 0) = x, \tag{108}$$

$$u_t(x, 0) = 1. \tag{109}$$

We have:

$$u_{tt} = c^2 u_{xx} - \alpha \sin u, \tag{110}$$

$$u_{tt}(x, 0) = c^2 u_{xx}(x, 0) - \alpha \sin u(x, 0), \tag{111}$$

$$u_{xx}(x, 0) = 0, \tag{112}$$

$$u_{tt}(x, 0) = -\alpha \sin x, \tag{113}$$

On equation (110), finding 1-order partial derivative of t on both sides, We have:

$$\tag{114}$$

$$u_{ttt}(x, 0) = c^2 u_{txx}(x, 0) - \alpha u_t(x, 0) \cos u(x, 0), \tag{115}$$

$$u_{txx}(x, 0) = 0, \tag{116}$$

$$u_{ttt}(x, 0) = -\alpha \cos x, \tag{117}$$

On equation (114), finding 1-order partial derivative of t on both sides, We have:

$$utttt = c2utttxx - autt \cos u + \alpha u^2 t \sin u, \quad (118)$$

$$u_{ttt}(x,0) = c^2 u_{tttx}(x,0) - \alpha u_{tt}(x,0) \cos u(x,0) + \alpha (u_t(x,0))^2 \sin u(x,0), \quad (119)$$

$$u_{tttx}(x,0) = \alpha \sin x, \quad (120)$$

$$u_{ttt}(x,0) = \alpha c^2 \sin x + \alpha^2 \sin x \cos x + \alpha \sin x,$$

$u_{tttt}(x,0), u_{ttttt}(x,0) \dots$  is known function.

By Taylor's formula, we get as following:

$$u(x,t) = u(x,0) + u_t(x,0)t + u_{tt}(x,0)\frac{t^2}{2!} + u_{ttt}(x,0)\frac{t^3}{3!} + u_{tttt}(x,0)\frac{t^4}{4!} + \dots \quad (121)$$

So, we can get the solution of (107).

Iterative solution of partial differential equations by Taylor formula is important and good method that solve linear and nonlinear partial differential equations. And the method also can solve fractional partial differential equations.

## References

1. A Bahri and H.Br'ézis (1980) periodic solution of a nonlinear wave equation, Proc. ROY.SOC.edinburgh Sect. A85.
2. A Bamberger, G Chavent and P Lailly (1979) About the stability of the inverse problem in 1-D wave equations. Applications to the interpretation of seismic profiles, Appl.Math.Optim 5: 1-47.
3. V Barbu and N H Pavel (1996) Periodic solutions to one-dimensional wave equation with piece-wise constant coefficient, J.Differential Equations 132: 319-337.
4. V Barbu and N H Pavel (1997) Determining the acoustic impedance in the 1-D wave equation via an optimal control problem, SIAMJ. Control Optim 35: 1544-1556.
5. V Barbu and N H Pavel (1997) Periodic solution to nonlinear one-dimensional wave equation with x-dependent coefficients, Trans.Amer.MATH.SOC 349: 2035-2048.
6. H Br'ézis (1983) Periodic solutions of nonlinear vibrating strings and duality principles, Bull.Amer.Math. Soc.(N.S.) 8: 409-426.
7. H Br'ézis and L Nirenberg (1978) Forced vibrations for a nonlinear wave equation, Comm.Pure Appl. Math 31: 1-30.
8. W Craig and CE Wayne (1993) Newton's method and periodic solutions of nonlinear wave equations, Cmm. pure Appl.Math 46: 1409-1498.
9. Y Ding S Li and M willem (1998) P eriodic Solutions Of Symmetric wave equations, J.Differential Equations 145: 217-241.
10. Y Gao, HeYuan Wang (2015) Taylor's formula and partial differential equation, Studies in College Mathematics 01-0079-03, 1008-1399.